

This is a repository copy of *The natural diversity and ecology of fission yeast*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/124871/>

Version: Accepted Version

Article:

Jeffares, Daniel C orcid.org/0000-0001-7320-0706 (2018) The natural diversity and ecology of fission yeast. *Yeast* (Chichester, England). pp. 1-21. ISSN 0749-503X

<https://doi.org/10.1002/yea.3293>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1 **Title:** The natural diversity and ecology of fission yeast.

2 **Running head:** Natural fission yeast diversity and ecology

3

4 Daniel C. Jeffares

5 Department of Biology

6 University of York

7 Wentworth Way

8 York YO10 5DD

9 United Kingdom

10 Email: daniel.jeffares@york.ac.uk

11

12 Keywords: *Schizosaccharomyces*, genetic diversity, fission yeast

13

14

15

16 **Abstract**

17 While the fission yeast is a powerful model of eukaryote biology, there have been
18 few studies of quantitative genetics, phenotypic or genetic diversity. Here I survey
19 the small collection of fission yeast diversity research. I discuss what we can infer
20 about the ecology and origins of *Schizosaccharomyces pombe* from microbiology
21 field studies and the few strains that have been collected.

22

23 **Introduction**

24 *Schizosaccharomyces pombe* research began in the 1940s (Fantes and Hoffman
25 2016) and is now a potent model of eukaryote biology, with a well-annotated
26 curated genome (Wood *et al.* 2002; McDowall *et al.* 2015), an extensive battery
27 of technical methods and genome-scale tools (Hoffman, Wood and Fantes 2015;
28 Hagan *et al.* 2016) and regular international meetings devoted to its study. Part of
29 the important utility of fission yeast as a model is that it contains many vertebrate
30 orthologs that are not present in budding yeast (Hoffman, Wood and Fantes
31 2015), so it provides a complement for studies of cell biology.

32 The majority of fission yeast research has used the strains described by
33 Leupold with its three mating types (Leupold 1949), and mutants derived from
34 these strains. Studies of diversity or quantitative genetics have been few and far
35 between. By contrast there is an extensive literature describing diversity and
36 quantitative genetics in the budding yeast *Saccharomyces cerevisiae* and its wild
37 relative *Saccharomyces paradoxus*, and a range of related species (Peter and
38 Schacherer 2016). These include QTL studies

39 (Swinnen, Thevelein and Nevoigt 2012; Liti and Louis 2012; Fay 2013;
40 Bloom *et al.* 2013; Märtens *et al.* 2016), genome-scale analysis of diversity (Liti
41 *et al.* 2009; Schacherer *et al.* 2009) and analysis of diversity and evolution in the
42 natural environment (Robinson, Pinharanda and Bensasson 2016; Leducq *et al.*
43 2016). In this review, I survey fission yeast diversity research, and I discuss what
44 little is known about the origins and natural ecology of this species.

45

46 **Defining fission yeast species**

47 Collections of *Schizosaccharomyces* strains were classified into three groups based
48 on crossing and protoplast fusion (Sipiczki *et al.* 1982), phenotypic characters
49 (Bridge and May 1984), DNA optical reassociation and physiological characters

50 (Vaughan Martini 1991), simplifying the rather complex list of potential ‘species’
51 into three (*Schizosaccharomyces pombe*, *S. japonicus*, *S. octosporus*).
52 *Schizosaccharomyces cryophilus* was identified much later as a contaminant of a
53 *S. octosporus* strain (CBS7191) from Denmark, and the species description was
54 accompanied by a draft genome (Helston *et al.* 2010).

55 The genomes and transcriptomes of *S. japonicus*, *S. octosporus* and an
56 improved *S. cryophilus* genome were described in 2011, showing that the
57 *Schizosaccharomyces* genus is as divergent on the protein level as the human-
58 amphioxus divergence (~55% amino acid identity) (Rhind *et al.* 2011). This
59 analysis described the conservation of orthologous groups, conservation of
60 transcription, the evolution of mating type regions and transposons. It also
61 features the first sequencing of a non-reference strain of *S. pombe*, concluding that
62 the within-species diversity was < 1% (confirmed later with studies of more
63 strains (Fawcett *et al.* 2014; Jeffares *et al.* 2015)). The current clade of only four
64 highly divergent fission yeast species is a limitation for evolutionary studies, since
65 evolutionary constraints can be estimated only inaccurately, and non-coding sites
66 that are in general subject to weaker purifying selection tend to be saturated
67 (Rhind *et al.* 2011). None of the *Schizosaccharomyces* species is sufficiently
68 closely related to *S. pombe* to reliably determine ancestral nucleotide states.

69

70 **Early (pre-genome sequence) diversity studies**

71 An early field study of this species was conducted by Florenzano *et al.*, who
72 showed that *S. pombe* was frequently present on grapes in Sicilian vineyards
73 (Florenzano, Balloni and Materassi 1977). Phenotypic characterization began with
74 analysis of xerotolerance (resistance to high solute concentrations) in 27 *S. pombe*
75 strains (Ganthala, Marshall and May 1994). One the first genetic analysis of
76 diversity within *S. pombe* described the intron content of mitochondrial genomes
77 in 26 strains, showing presence/absence polymorphisms in group I and group II
78 introns (Zimmer *et al.* 1987). Interestingly, there appears to be no intron presence
79 polymorphisms in the nuclear genomes of sequenced strains (Mourier & Jeffares,
80 unpublished analyses), though on the longer scale fission yeasts have certainly
81 undergone intron gain and loss (Mourier and Jeffares 2003; Jeffares, Mourier and
82 Penny 2006; Rhind *et al.* 2011).

83 In a prelude to genome-scale analyses, three studies began to explore
84 genetic and phenotypic diversity on a larger scale. Gomes *et al.*, collected 27

85 strains from seven Brazilian cachaça distilleries, and characterised osmotolerance,
86 trehalose accumulation and ethanol tolerance, showing that these strains could
87 grow in 50% glucose and 10% ethanol (Gomes *et al.* 2002). They also explored
88 population structure using RAPD-PCR (random amplified polymorphic DNA
89 PCR), demonstrating local population structure in Brazilian cachaça strains.
90 RAPD-PCR was a useful method to characterise diversity prior to next generation
91 sequencing, but the development of 26 primers for microsatellite PCR now
92 provide a simple method to genotype strain collections (Patch and Aves 2007).
93 Brown *et al.* assembled 81 natural isolates of *S. pombe* including samples from all
94 continents (except Antarctica), and measured a large assembly of phenotypic
95 characters, including growth parameters in 42 liquid media and cell length (Brown
96 *et al.* 2011). This analysis also described diversity at three locations, and
97 estimated that the global effective population size of this species is 10^7 (a figure
98 that remained after genome-wide analysis (Farlow *et al.* 2015)). Most
99 interestingly, this work described extensive karyotype diversity within this
100 collection, including reciprocal translocations, duplications and inversions,
101 showing that the ribosomal repeats were located on different chromosome ends in
102 different strains.

103

104 **Genome-wide sequence analyses**

105 The creation and analysis of the only fission yeast recombinant strain library
106 was published in 2014 (Clément-Ziza *et al.* 2014). This study used a two-parent
107 segregant panel and described expression QTLs (eQTLs) from both protein-
108 coding and non-coding transcripts, during growth and stress conditions.
109 Interestingly this study discovered a larger proportion of associations between
110 genetic variants and non-coding transcripts than coding transcripts. The most
111 significant variant, that affected 44% of eQTL associations and growth rate, was a
112 frameshift in the *swc5* gene - part of a complex that affects histone deposition.
113 Detailed analysis showed that this frameshift caused increased antisense
114 transcription and decreased sense transcription, providing an example of the
115 molecular events that influenced a complex trait such as growth. Further analyses
116 of segregant panels are in progress, describing positive selection and the genetic
117 control of RNA and protein levels (Clément-Ziza, pers. comm.).

118 An analysis of segregant pool based mapping (bulk segregant analysis) from
119 a two-parent cross showed that this method was feasible in fission yeast (Hu, Suo

120 and Du 2015). Hu *et al.* localised the probable causal allele of maltose deficiency
121 by sequencing pools grown with and without maltose. The analysis was
122 complicated by an inversion in the reference strain, but few other wild strains
123 (Jeffares *et al.* 2017), which reduces the local recombination rate (Clément-Ziza *et*
124 *al.* 2014).

125 Two genome-wide analyses of genetic diversity in *S. pombe* were published
126 soon afterwards (Fawcett *et al.* 2014; Jeffares *et al.* 2015). Both analyses
127 described recombination rate and population structure, and showed that exons,
128 UTRs and introns were the main targets of purifying selection. Estimates of
129 diversity (π) were $\sim 3 \times 10^{-3}$ (pairwise comparison have an average of 3 SNPs/kb),
130 slightly higher than the budding yeast *Saccharomyces cerevisiae* (1×10^{-3}) (Liti *et*
131 *al.* 2009). From the genetic diversity and mutation rates, the effective population
132 size of *S. pombe* has been estimated to be 12 million, on a similar scale to budding
133 yeast (3 million) (Farlow *et al.* 2015).

134 The analysis of Fawcett *et al.* (32 strains) described some unusual patterns
135 of diversity that were likely due to soft selective sweeps, and either balancing
136 selection or introgression from some unknown fission yeast outgroup (Fawcett *et*
137 *al.* 2014). Jeffares *et al.* (161 strains) described transposon insertions and included
138 analysis of quantitative traits, their heritability and quantitative genetics using the
139 genome-wide association study (GWAS) approach (Jeffares *et al.* 2015). This
140 study located 1,400 variants that were significantly associated with traits despite
141 the very small sample size, showing that the combination of simple tractable
142 genetics with the capability to measure traits accurately with abundant repeat
143 measurements in well-controlled environments, is a powerful combination.
144 Further analysis with the same strain collection described structural variants
145 showing that they are both transient and contribute considerably to quantitative
146 traits and reproductive isolation (Jeffares *et al.* 2017). Interestingly the variance in
147 wine-making traits, such as malic acid accumulation and glucose/fructose
148 utilisation (Benito *et al.* 2016), appeared to be caused entirely by structural
149 variants.

150 Two genome-scale analyses of the mutation rate estimated the point
151 mutation rate to be 1.7×10^{-10} (or 2.0×10^{-10}) per base per generation (Farlow *et*
152 *al.* 2015; Behringer and Hall 2015), very similar to estimates for the budding
153 yeast *Saccharomyces cerevisiae* (estimates at 3 and 1.67×10^{-10}) (Lynch *et al.*
154 2008; Zhu *et al.* 2014). Both studies noted a strong bias towards small insertions,

155 over deletions, which occur primarily in the non-protein regions of the genome, a
156 pattern that is retained in natural genetic diversity (Jeffares *et al.* 2015).

157

158 **Reproductive isolation**

159 One topic that has received particular attention is the study of mating types
160 and reproductive isolation. Since the outset of fission yeast research, it was clear
161 homothallic strains could mutate to more or less stable heterothallic genotypes (h^+
162 or h^-) (Leupold 1949). Natural isolates also vary genetically at mating type
163 regions and in their mating behavior, with some strains mutating more frequently
164 from h^+ to h^- and vice versa (Schlake and Gutz 1993). In an interesting
165 demonstration that reproductive isolation could evolve via pre-zygotic
166 mechanisms, Sieke *et al.* created three novel reproductive groups with different
167 pheromone-receptor pairs (Seike, Nakamura and Shimoda 2015). Given these
168 changes it is likely that pre-zygotic reproductive isolation occurs within some
169 populations.

170 Several studies described the low spore viability that results from many
171 inter-strain matings (Kondrat'eva and Naumov 2001; Teresa Avelar *et al.* 2013;
172 Zanders *et al.* 2014; Naumov and Kondratieva 2015; Jeffares *et al.* 2015).
173 Viability ranges from pairs showing $< 1\%$ viable offspring to strains with 90%
174 viable, similar a range observed for *species* of budding yeast with that have much
175 higher genetic divergence than fission yeast strains (Liti, Barton and Louis 2006),
176 consistent with *S. pombe* strains being 'on the verge of speciation' (Naumov and
177 Kondratieva 2015) (**Figure 1A**). Some homothallic strains are also ineffective at
178 mating with their own genotype (Kondrat'eva and Naumov 2001; Jeffares *et al.*
179 2015).

180 Since most crosses do produce mating bodies and asci (Xavi Marsellach,
181 pers. comm.), the isolation is generally post-zygotic (intrinsic reproductive
182 isolation). The accumulation of genetic factors that reduce mating success
183 within these relatively closely related strains is probably due to the low
184 frequency of outbreeding in fission yeast. Based on the decay in linkage between
185 wild isolates Farlow *et al.* estimated that *S. pombe* mate with a genetically
186 dissimilar individual on average every 800,000 generations (Farlow *et al.* 2015),
187 far less frequently than the estimates 50,000 generation for *S. cerevisiae* (Ruderfer
188 *et al.* 2006). Given this frequency of, it is not surprising that the existing strains

189 have accumulated genetic factors that preclude interbreeding in the ~2300 years
190 since these strains have drifted apart (Jeffares *et al.* 2015).

191 There are at least three (non-exclusive) genetic causes for the reproductive
192 isolation of fission yeasts. Spore killing (meiotic drive), has been proposed to be a
193 mechanism (Kondrat'eva and Naumov 2001; Zanders *et al.* 2014; Naumov and
194 Kondratieva 2015). Many of the crosses analysed by Kondratieva *et al.* from
195 genetically divergent strains and produced strong deviations from expected
196 Mendelian ratios (Kondrat'eva and Naumov 2001; Naumov, Kondratieva and
197 Naumova 2015) (Figure 1B), while the analyses of Zanders *et al.* concluded that
198 there were meiotic drive elements on all three chromosomes (Zanders *et al.* 2014).

199 Two recent analyses have demonstrated that members of the *wtf* gene
200 family mediate drive with a spore killer-antidote system (Hu *et al.* 2017; Nuckolls
201 *et al.* 2017). Hu *et al.* demonstrate that *wtf9* and *wtf27* genes from the non-
202 reference strain (CBS5557/JB4) drive segregation distortion in when mated to the
203 reference strain, that this drive is independent of genomic location. Nuckolls *et al.*
204 show that *wtf4* promotes distortion in crosses between the reference strain and the
205 kombucha strain (SPK1820/YFS276/JB1180, as initially sequenced by the Broad
206 Institute (Rhind *et al.* 2011)). Other strains analysed by Kondratieva *et al.* also
207 show very biased segregation (Figure 1B).

208 Collectively, these analyses show that the spore killer (or poison) and
209 antidote functions can be separated by mutations. In the natural state, there are
210 two transcripts that mediate killer/antidote functions (Nuckolls *et al.* 2017). While
211 the killer protein variant is distributed in all four spores of the asci, the antidote
212 remains only within cells with the relevant *wtf* genotype. Since *wtf* genes encode
213 membrane-spanning domains they may travel between asci. The genetics of the
214 poison-antidote systems are complex, in that there are multiple *wtf* genes in
215 different strains that have degenerated to contain the poison and antidote
216 functions, antidote only, or no function. Both analyses show that *wtf* genes are
217 particularly genetically diverse (Figure 1C). However, they do not show an excess
218 of high Tajima's D values (Tajima 1989)(Figure 1C), a genetic diversity
219 parameter which is one of the expected signatures of balancing selection.

220 Reproductive isolation may also be the result of the aneuploidy that occurs
221 when parents differ in chromosomal inversions and translocations. For example,
222 engineered inversions and translocations reduce spore viability by ~40% (Teresa
223 Avelar *et al.* 2013). *S. pombe* strains do have extensive karyotype differences

224 (Brown *et al.* 2011; Naumov, Kondratieva and Naumova 2015; Jeffares *et al.*
225 2017), including a strain that maintains four (rather than the usual three)
226 chromosomes (Brown *et al.* 2014). There is a significant association between
227 viability and the SV-distance between parents (Jeffares *et al.* 2017), though
228 viability declines at less than 40% viability per variant. This is probably because
229 natural structural variants are biased to chromosome ends that do not contain
230 essential genes (Jeffares *et al.* 2015), due to selection for those that do not cause
231 lethal aneuploidies. Structural variants may also contribute to drive (Zanders *et al.*
232 2014).

233 Formally, reproductive isolation may also be due to Bateson-Dobzhansky-
234 Muller interactions (BDMIs) or any of the other genetic mechanisms of negative
235 epistasis (Nei and Nozawa 2011). However segregation data from random spores
236 (Kondrat'eva and Naumov 2001; Naumov and Kondratieva 2015) and dissected
237 tetrads is inconsistent with simple two-locus BDMIs, which are expected to
238 produce small deviations from expected segregation patterns (even when the
239 affected alleles were strongly linked to markers) (Hou and Schacherer 2016).
240 Ultimately meiotic drive, epistasis and structural variants may have interacting
241 effects on viability, since locally adapted haplotypes are predicted to develop
242 within areas of reduced recombination (Kirkpatrick and Barton 2006).

243 With all these studies of population genetics (reproductive isolation,
244 divergence dating, diversity measures, population size *etc.*) the analyses are based
245 on a small collection of strains that are a worldwide sample of mostly human
246 commensals (see below), so conclusions may not represent natural populations.
247




Figure 1. Intrinsic reproductive isolation in *S. pombe*.

A) Random spore viability from three studies shows a decline in spore survival with genetic distance (SNP distance) between parents. The size of circles indicates the lowest self-mating viability of parents. Data from (Kondrat'eva and Naumov 2001; Teresa Avelar *et al.* 2013; Jeffares *et al.* 2015). Crosses involving the strain CBS5680 (as in part B) are indicated with cross hairs. The range of genetic differences that have highly variable effects on viability (10,000 – 30,000 SNPs) is indicated by vertical dashed lines. The outlier at top right is JB848/CBS10475 (Brazil) x JB870/CBS10499 (South Africa), which appears to be real (Xavier Marsellach, pers. comm.). **B)** segregation of control markers in random spore analysis show strong deviations from the expected 1:1:1:1 ratio, data from (Kondrat'eva and Naumov 2001). For one strain (CBS5680/JB873, from Poland) we show the counts of control markers (aB and Ab are parental types, AB, ab are recombinants, see Kondrat'eva *et al.* for details). Segregation counts whose χ^2 test P-values were < 0.05 are plotted with red bars. Plot text shows the parents of the cross, the random spore viability (RSV) and the χ^2 test P-value (CHISQ.P). **C)** *wtf* genes have high pairwise diversity within strains compared to all other transmembrane domain containing and non-TM genes (π , left panel), high numbers of segregating sites (θ , middle panel), but are not outliers for Tajima's D (which is calculated from the ratio of the two, D, right panel). Plots show diversity estimators from 57 strains, red circle indicate individual values for *wtf* genes.

271 Predicted transmembrane proteins were collected from a query of Pombase
272 (www.pombase.org), diversity data from (Jeffares *et al.* 2015).

273

274 **Genetics and the reference strain**

275 The fission yeast community has worked almost exclusively with one reference
276 strain, and spontaneous mutants generated from this strain (Fantes and Hoffman
277 2016). This laboratory strain is a natural isolate, and is not an unusual strain
278 phenotypically. It does not appear to be adapted to the standard rich or minimal
279 media, since it does not grow particularly rapidly in these media compared to wild
280 strains. There are several important discoveries that are relevant to the fission
281 yeast researcher. Firstly, Wild strains can differ from the reference by up to
282 68,000 SNPs and up to 24 structural variations, which contribute to phenotypic
283 variation between strains (Clément-Ziza *et al.* 2014; Jeffares *et al.* 2015; Hu, Suo
284 and Du 2015; Jeffares *et al.* 2017). I summarise the structural differences between
285 strains in Supplementary Figure 1. Secondly, the structural differences and
286 meiotic drive elements that wild strains contain complicate crosses between
287 strains, by reducing spore viability and skewing the proportions of alleles that are
288 produced in the offspring (Kondrat'eva and Naumov 2001; Kondrateva and
289 Naumov 2011; Clément-Ziza *et al.* 2014; Hu, Suo and Du 2015; Nuckolls *et al.*
290 2017; Hu *et al.* 2017).

291

292 **The ecology of fission yeast**

293 There have been few published attempts to systematically collect fission
294 yeast strains (Gomes *et al.* 2002; Benito *et al.* 2013; Hellberg 2013). However,
295 fission yeasts have been serendipitously discovered in a variety of microbiological
296 studies (Table 1, Figure 2). Sources have generally been traditional non-
297 industrialised fermentations, produced without any intentional inoculation from
298 substrates that contain high concentrations of sugars. When quantitative estimates
299 of species abundances are included *Schizosaccharomyces* yeasts were generally
300 minor components of these fermentations, with the exceptions of kombucha, some
301 cachaça fermentations and baijiu (from tea, sugar cane and sorghum respectively)
302 (Pataro, Guerra and Peixoto 2000; Teoh, Heard and Cox 2004; Wu, Xu and Chen
303 2012).

304 Perhaps more informative for fission yeast ecology, are the cases where

305 fission yeasts have been discovered in natural substrates such as palm wine (a
 306 fermentation of palm sap) (Theivendirarajah and Chrystopher 1987;
 307 Amanchukwu, Obafemi and Okpokwasili 1989; Ouoba *et al.* 2012). Fission yeast
 308 are also present in natural fermentations of fruits such as *Coffea arabica* and
 309 *Theobroma cacao* (from which coffee and cocoa beans are harvested respectively)
 310 (Silv *et al.* 2000; Schwan and Wheals 2004). Collectively, the field studies show
 311 that fission yeasts are a component of natural microbial communities that ferment
 312 botanical sugars in several geographic regions.

313 Including the strains present in stock collections and in field studies the
 314 most common substrates for fission yeast have been palm wine, grape wine, high-
 315 sugar substrates (molasses, cane sugar, honey) and fruits (Figure 2). Three
 316 selective media to have been described to enrich for fission yeast (Florenzano,
 317 Balloni and Materassi 1977; Hellberg 2013; Benito *et al.* 2013), so further
 318 systematic collections from similar locations and substrates should be possible in
 319 the future.

320

321 **Table 1. *Schizosaccharomyces* in field microbiology**

Substrate	Location	Reference
Grape must	Sicily	(Florenzano, Balloni and Materassi 1977)
Grapes	Ukraine	(Bayraktar 2014)
Palm wine	Sri Lanka	(Atputharajah, Widanapathirana and Samarajeewa 1986; Theivendirarajah and Chrystopher 1987)
Palm wine	Nigeria	(Sanni and Lönner 1993; Amanchukwu, Obafemi and Okpokwasili 2006)
Palm wine	Burkina Faso	(Ouoba <i>et al.</i> 2012)
Rum	Haiti	(Fahrasmane, Ganou-Parfait and Parfait 1988)
Molasses, raisin	Japan/Thailand/Taiwan	(Ishitane 1985)


Tequila	Mexico	(Lachance 1995)
Coffee cherries	Brazil Madagascar	(Silv <i>et al.</i> 2000) (Ravelomanana <i>et al.</i> 1984)
Cachaça (from sugar cane)	Brazil	(Pataro, Guerra and Peixoto 2000; Gomes <i>et al.</i> 2002)
Kombucha (fermented tea)	Australia**	(Teoh, Heard and Cox 2004)
Cocoa pulp	Belize	(Schwan and Wheals 2004)
Baijiu (distillate of fermented sorghum)	China	(Wu, Xu and Chen 2012)
Traditional breweries	China	Fen-Yang Bai, pers. comm.
Honey	Fiji	(Ponici and Wimmer 1986)
Honey	Spain	(Benito <i>et al.</i> 2014)

322 * Not microbiological study itself, refers to earlier work.

323 ** From commercial kombucha brewers.

324

325



326

327

328 **Figure 2. Fission yeast locations and substrates.** The locations and
329 substrates where fission yeast have been discovered, including all strains that have
330 been sequenced from stock centers (Fawcett *et al.* 2014; Jeffares *et al.* 2015), and
331 reports from field studies (Table 1). Sequenced strains are marked with cross-
332 hairs, and strains isolated from uncertain locations are marked with a square.

333

334 **The origin of fission yeast**

335 *S. pombe* is now globally distributed (Figure 2), but we know little about its
336 origin and dispersal. We have estimated that these strains began to spread globally
337 in from ~340 BCE (95% confidence interval 1875 BCE–1088 CE), and that the
338 current collection of strains from Brazilian cachaça originated from the remainder
339 in about ~1620 CE (confidence interval 1422–1752 CE) (Jeffares *et al.* 2015), a
340 hint that like budding yeast and *C. elegans*, this model has probably been
341 dispersed as a commensal (most likely in fermented beverages).

342 The reference strain originated from French grapes (Osterwalder 1924). The
343 common belief is that *S. pombe* originated from Africa, perhaps because the initial
344 species description was from an African millet beer isolate (Lindner 1893;
345 Vorderman 1894). While genetic analysis is consistent with exchange between
346 African and European stocks (Jeffares *et al.* 2015), and some strains have been
347 collected from traditional African fermentations, there is no scientific evidence for
348 an African origin of this species. There are very few studies of the microbial
349 constituents of millet beer from Africa (I could find none than specifically
350 mentioned *S. pombe*, and one description of sorghum beer that did not mention *S.*
351 *pombe* (Kayode *et al.* 2011)). Since fission yeasts can be major components of
352 kombucha, which has been traditionally produced in China (Sreeramulu, Zhu and
353 Knol 2000; Teoh, Heard and Cox 2004), palm wine which is widely produced in
354 Asia (Table 1, Figure 2), and in traditional Chinese breweries (Fen-Yang Bai,
355 pers. comm.), China is an equally good candidate for the initial origin of *S.*
356 *pombe*.

357

358 **Why study diversity in fission yeast?**

359 The small genomes of budding yeasts enabled the early development of
360 population genomics methods (Liti *et al.* 2009; Schacherer *et al.* 2009), and now
361 large scale accurate quantitative genetics analyses (Bloom *et al.* 2013; Märtens *et*
362 *al.* 2016). The continuing advance of sequence throughput, analysis software and

laboratory methods (eg: RAD-seq) have now made population genomics approaches available to any species. However, the abundance of genome-scale data and technical tools and the small non-redundant genomes of yeasts make them attractive models for systems biology, including approaches to understanding genetic diversity and traits (Parts 2014). Fission yeast has the benefit of being haploid (so that F1 generations need not be intercrossed). As with budding yeast, fission yeast has abundant heritable phenotypic diversity in growth, stress responses, cell morphology, and cellular biochemistry that is yet to be explored with powerful quantitative genetics (Brown *et al.* 2011; Clément-Ziza *et al.* 2014; Jeffares *et al.* 2015; 2017). Yeasts are also powerful tools for detailed study of evolutionary processes using pooled time-series sequencing and other high-throughput approaches that would be expensive or unfeasible in other species (Cubillos *et al.* 2011; Hou *et al.* 2015). Finally, studies by Benito *et al.* show that some non-reference *S. pombe* strains have potential in the winemaking industry (Benito *et al.* 2014; 2016), so diverse strains could well have potential elsewhere in biotechnology.

379

380 **Acknowledgements**

381 I thank Mathieu Clément-Ziza for commentary about unpublished work and
382 Xavier Marsellach for discussions.

383

384 **Supplementary data**

385 All used for plots is available at figshare at:

386 https://figshare.com/projects/The_natural_diversity_and_ecology_of_fission_yeast/21761
387 [t /21761](https://figshare.com/projects/The_natural_diversity_and_ecology_of_fission_yeast/21761)

388

389

390

391

392


393

394

395

396

397



399

400

401 **Supplementary Figure 1. Structural variants present in wild fission yeast**
402 **strains.** Using predictions from short read data (Jeffares *et al.* 2017), I show the
403 genomic location of structural variants (SVs) in wild strains contain that differ
404 from the standard laboratory isolate (Leupold’s 972) . I show deletions (black),
405 duplications (red), inversions (green) and translocations (blue). SVs present in
406 each of the 57 non-clonal strains are shown within the white horizontal bars, with
407 strain names coloured according to their continent of origin. Tf1-type
408 retrotransposon insertions that are present in some, but not all strains are shown at
409 grey ticks at the tops of bars. The positions of fixed Tf1-type retrotransposon
410 insertions are indicated on the last row (f/LTRs). Centromeres are indicated with
411 black triangles.

412

413

414

415 **References**

416

417 Amanchukwu SC, Obafemi A, Okpokwasili GC. Single-cell-protein production
418 by *Schizosaccharomyces pombe* isolated from palmwine using hydrocarbon
419 feedstocks. *Folia Microbiol* 1989;**34**:112–9.

- 420 Amanchukwu SC, Obafemi A, Okpokwasili GC. Hydrocarbon degradation and
421 utilization by a palm-wine yeast isolate. *FEMS microbiology letters*
422 2006;**57**:151–4.
- 423 Atputharajah JD, Widanapathirana S, Samarajeewa U. Microbiology and
424 biochemistry of natural fermentation of coconut palm sap - ScienceDirect.
425 *Food Microbiol* 1986.
- 426 Bayraktar VN. Biodiversity of yeast cultures isolated from littoral areas of the
427 Regional Landscape Park «Tiligulskiy». *J. Optimization and Protection of*
428 *Ecosystems* 2014.
- 429 Behringer MG, Hall DW. Genome-Wide Estimates of Mutation Rates and
430 Spectrum in *Schizosaccharomyces pombe* Indicate CpG Sites are Highly
431 Mutagenic Despite the Absence of DNA Methylation. *G3* 2015;**6**:149–60.
- 432 Benito Á, Jeffares DC, Palomero F *et al.* Selected *Schizosaccharomyces pombe*
433 Strains Have Characteristics That Are Beneficial for Winemaking. *PLoS ONE*
434 2016;**11**:e0151102.
- 435 Benito S, Gálvez L, Palmero F *et al.* *Schizosaccharomyces* selective differential
436 media. *African Journal of Microbiology Research* 2013;**7**:3026–36.
- 437 Benito S, Palomero F, Calderón F *et al.* Selection of appropriate
438 *Schizosaccharomyces* strains for winemaking. *Food Microbiol* 2014;**42**:218–
439 24.
- 440 Bloom JS, Ehrenreich IM, Loo WT *et al.* Finding the sources of missing
441 heritability in a yeast cross. *Nature* 2013;**494**:234–7.
- 442 Bridge PD, May JW. A numerical classification of fission yeasts of the genus
443 *Schizosaccharomyces* Lindner. *Microbiology* 1984;**130**:1921–32.
- 444 Brown WRA, Liti G, Rosa C *et al.* A geographically diverse collection of
445 *Schizosaccharomyces pombe* isolates shows limited phenotypic variation but
446 extensive karyotypic diversity. Fay JC (ed.). *G3* 2011;**1**:615–26.
- 447 Brown WRA, Thomas G, Lee NCO *et al.* Kinetochore assembly and
448 heterochromatin formation occur autonomously in *Schizosaccharomyces*

449 pombe. *Proc Natl Acad Sci USA* 2014;**111**:1903–8.

450 Clément-Ziza M, Marsellach FX, Codlin S *et al.* Natural genetic variation impacts
 451 expression levels of coding, non-coding, and antisense transcripts in fission
 452 yeast. *Mol Syst Biol* 2014;**10**:764–4.

453 Cubillos FA, Billi E, Zörgö E *et al.* Assessing the complex architecture of
 454 polygenic traits in diverged yeast populations. *Mol Ecol* 2011;**20**:1401–13.

455 Fahrasmane L, Ganou-Parfait B, Parfait A. Research note: Yeast flora of Haitian
 456 rum distilleries. *Mircen Journal* 1988;**4**:239–41.

457 Fantes PA, Hoffman CS. A Brief History of *Schizosaccharomyces pombe*
 458 Research: A Perspective Over the Past 70 Years. *Genetics* 2016;**203**:621–9.

459 Farlow A, Long H, Arnoux S *et al.* The Spontaneous Mutation Rate in the Fission
 460 Yeast *Schizosaccharomyces pombe*. *Genetics* 2015;**201**:737–44.

461 Fawcett JA, Iida T, Takuno S *et al.* Population Genomics of the Fission Yeast
 462 *Schizosaccharomyces pombe*. *PLoS ONE* 2014;**9**:e104241.

463 Fay JC. The molecular basis of phenotypic variation in yeast. *Current Opinion in*
 464 *Genetics and Development* 2013;**23**:672–7.

465 Florenzano G, Balloni W, Materassi R. Contributo alla ecologia dei lieviti
 466 *Schizosaccharomyces* sulle uve. *Vitis* 1977;**16**:38–44.

467 Ganthala BP, Marshall JH, May JW. Xerotolerance in fission yeasts and the role
 468 of glycerol as compatible solute. *Archives of microbiology* 1994;**162**:108–13.

469 Gomes FCO, Pataro C, Guerra JB *et al.* Physiological diversity and trehalose
 470 accumulation in *Schizosaccharomyces pombe* strains isolated from
 471 spontaneous fermentations during the production of the artisanal Brazilian
 472 cachaça. *Can J Microbiol* 2002;**48**:399–406.

473 Hagan IM, Grallert A, Carr AM *et al.* eds. *Fission Yeast: a Laboratory Manual*.
 474 Cold Spring Harbour Laboratory Press, 2016.

475 Hellberg JEEU. Finding wild fission yeast: Where are they? *Genetics Society*
 476 *News* 2013;**68**:59–60.

- 477 Helston RM, Box JA, Tang W *et al.* Schizosaccharomyces cryophilus sp nov., a
478 new species of fission yeast. *Fems Yeast Res* 2010;**10**:779–86.
- 479 Hoffman CS, Wood V, Fantes PA. An Ancient Yeast for Young Geneticists: A
480 Primer on the Schizosaccharomyces pombe Model System. *Genetics*
481 2015;**201**:403–23.
- 482 Hou J, Friedrich A, Gounot J-S *et al.* Comprehensive survey of condition-specific
483 reproductive isolation reveals genetic incompatibility in yeast. *Nat Commun*
484 2015;**6**:7214.
- 485 Hou J, Schacherer J. On the Mapping of Epistatic Genetic Interactions in Natural
486 Isolates: Combining Classical Genetics and Genomics. *Methods Mol Biol*
487 2016;**1361**:345–60.
- 488 Hu W, Jiang Z-D, Suo F *et al.* A large gene family in fission yeast encodes spore
489 killers that subvert Mendel's law. *eLife* 2017;**6**:3025.
- 490 Hu W, Suo F, Du L-L. Bulk segregant analysis reveals the genetic basis of a
491 natural trait variation in fission yeast. *Genome Biol Evol* 2015;**7**:3496–510.
- 492 Ishitane. Identification of yeasts isolated from high-sugar foods. Tokuoka K,
493 Ishitane T, Goto S, et al. (eds.). *J Gen Appl Microbiol* 1985;**31**:411–27.
- 494 Jeffares DC, Jolly C, Hoti M *et al.* Transient structural variations have strong
495 effects on quantitative traits and reproductive isolation in fission yeast. *Nat*
496 *Commun* 2017;**8**:14061.
- 497 Jeffares DC, Jolly C, Hoti M *et al.* Transient structural variations have strong
498 effects on quantitative traits and reproductive isolation in fission yeast.
499 *Multiple values selected* **8**:14061EP–.
- 500 Jeffares DC, Mourier T, Penny D. The biology of intron gain and loss. *Trends*
501 *Genet* 2006;**22**:16–22.
- 502 Jeffares DC, Rallis C, Rieux A *et al.* The genomic and phenotypic diversity of
503 Schizosaccharomyces pombe. *Nature Genetics* 2015;**47**:235–41.
- 504 Kayode APP, Vieira-Dalode G, Linnemann AR *et al.* Diversity of yeasts involved
505 in the fermentation of tchoukoutou, an opaque sorghum beer from Benin.

- 506 *African Journal of Microbiology Research* 2011;**5**:2737–42.
- 507 Kirkpatrick M, Barton N. Chromosome inversions, local adaptation and
508 speciation. *Genetics* 2006;**173**:419–34.
- 509 Kondrat'eva VI, Naumov GI. The Phenomenon of Spore Killing in
510 *Schizosaccharomyces pombe* Hybrids. *Doklady Biological Sciences* 2001.
- 511 Kondrateva VI, Naumov GI. POPULATION ANTAGONISM IN THE YEASTS
512 *SCHIZOSACCHAROMYCES POMBE*. *Ecological genetics* 2011;**9**:21–6.
- 513 Lachance MA. Yeast communities in a natural tequila fermentation. *Antonie Van*
514 *Leeuwenhoek* 1995;**68**:151–60.
- 515 Leducq J-B, Nielly-Thibault L, Charron G *et al.* Speciation driven by
516 hybridization and chromosomal plasticity in a wild yeast. *Nat Microbiol*
517 2016;**1**:15003.
- 518 Leupold U. Die Vererbung von Homothallie und Heterothallie bei
519 *Schizosaccharomyces pombe*. *Compt Rend Trav Lab Carlsberg Ser*
520 1949;**24**:381–480.
- 521 Lindner P. *Schizosaccharomyces pombe* n. sp., ein neuer Gährungserreger.
522 *Wochenschrift für Brauerei* 1893;**10**:1298–300.
- 523 Liti G, Barton DBH, Louis EJ. Sequence diversity, reproductive isolation and
524 species concepts in *Saccharomyces*. *Genetics* 2006;**174**:839–50.
- 525 Liti G, Carter DM, Moses AM *et al.* Population genomics of domestic and wild
526 yeasts. *Nature* 2009;**458**:337–41.
- 527 Liti G, Louis EJ. Advances in quantitative trait analysis in yeast. Fay JC (ed.).
528 *PLoS Genet* 2012;**8**:e1002912.
- 529 Lynch M, Sung W, Morris K *et al.* A genome-wide view of the spectrum of
530 spontaneous mutations in yeast. *Proc Natl Acad Sci USA* 2008;**105**:9272–7.
- 531 Märtens K, Hallin J, Warringer J *et al.* Predicting quantitative traits from genome
532 and phenome with near perfect accuracy. *Nat Commun* 2016;**7**:11512.

- 533 McDowall MD, Harris MA, Lock A *et al.* PomBase 2015: updates to the fission
534 yeast database. *Nucleic Acids Res* 2015;**43**:D656–61.
- 535 Mourier T, Jeffares DC. Eukaryotic intron loss. *Science (New York, NY)*
536 2003;**300**:1393–3.
- 537 Naumov G, Kondratieva V. Hybrid sterility of the yeast *Schizosaccharomyces*
538 *pombe*: Genetic genus and many species in statu nascendi? *Microbiology*
539 (00262617) 2015.
- 540 Naumov GI, Kondratieva VI, Naumova ES. [Hybrid Sterility of the Yeast
541 *Schizosaccharomyces pombe*: Genetic Genus and Many Species in statu
542 nascendi?]. *Mikrobiologiya* 2015;**84**:192–203.
- 543 Nei M, Nozawa M. Roles of mutation and selection in speciation: from Hugo de
544 Vries to the modern genomic era. *Genome Biol Evol* 2011;**3**:812–29.
- 545 Nuckolls NL, Bravo Núñez MA, Eickbush MT *et al.* wtf genes are prolific dual
546 poison-antidote meiotic drivers. *eLife* 2017;**6**:2235.
- 547 Osterwalder A. *Schizosaccharomyces liquefaciens* n.sp., eine gegen freie
548 schweflige Säure widerstandsfähige Gärhefe. *Mitt Geb Lebensmittelunters*
549 *Hyg* 1924;**15**:5–28.
- 550 Ouoba LII, Kando C, Parkouda C *et al.* The microbiology of Bandji, palm wine of
551 *Borassus akeassii* from Burkina Faso: identification and genotypic diversity
552 of yeasts, lactic acid and acetic acid bacteria. *J Appl Microbiol* 2012, DOI:
553 10.1111/jam.12014.
- 554 Parts L. Genome-wide mapping of cellular traits using yeast. *Yeast* 2014;**31**:197–
555 205.
- 556 Pataro C, Guerra JB, Peixoto MP. Yeast communities and genetic polymorphism
557 of *Saccharomyces cerevisiae* strains associated with artisanal fermentation in
558 Brazil. *Journal of Applied Microbiology* 2000;**89**:24–31.
- 559 Patch A-M, Aves SJ. Fingerprinting fission yeast: polymorphic markers for
560 molecular genetic analysis of *Schizosaccharomyces pombe* strains.
561 *Microbiology (Reading, Engl)* 2007;**153**:887–97.

- 562 Peter J, Schacherer J. Population genomics of yeasts: towards a comprehensive
563 view across a broad evolutionary scale. *Yeast* 2016;**33**:73–81.
- 564 Ponici L, Wimmer FL. Characterization of the yeasts (Blastomycetes) in some
565 Fijian honeys. *Acta alimentaria polonica* 1986.
- 566 Ravelomanana R, Guiraud JP, Vincent JC *et al.* Study of the yeast flora from the
567 traditional cacao fermentation in Madagascar. *Revue des Fermentations et des*
568 *Industries Alimentaires (Belgium)* 1984.
- 569 Rhind N, Chen Z, Yassour M *et al.* Comparative functional genomics of the
570 fission yeasts. *Science (New York, NY)* 2011;**332**:930–6.
- 571 Robinson HA, Pinharanda A, Bensasson D. Summer temperature can predict the
572 distribution of wild yeast populations. *Ecol Evol* 2016;**6**:1236–50.
- 573 Sanni AI, Lönner C. Identification of yeasts isolated from Nigerian traditional
574 alcoholic beverages. *Food Microbiol* 1993;**10**:517–23.
- 575 Schacherer J, Shapiro JA, Ruderfer DM *et al.* Comprehensive polymorphism
576 survey elucidates population structure of *Saccharomyces cerevisiae*. *Nature*
577 2009;**458**:342–5.
- 578 Schlake T, Gutz H. Mating configurations in *Schizosaccharomyces pombe* strains
579 of different geographical origins. *Curr Genet* 1993.
- 580 Schwan RF, Wheals AE. The microbiology of cocoa fermentation and its role in
581 chocolate quality. *Crit Rev Food Sci Nutr* 2004;**44**:205–21.
- 582 Seike T, Nakamura T, Shimoda C. Molecular coevolution of a sex pheromone and
583 its receptor triggers reproductive isolation in *Schizosaccharomyces pombe*.
584 *Proc Natl Acad Sci USA* 2015;**112**:4405–10.
- 585 Silv CF, Schwan RF, Sousa Dias ES *et al.* Microbial diversity during maturation
586 and natural processing of coffee cherries of *Coffea arabica* in Brazil. *Int J*
587 *Food Microbiol* 2000;**60**:251–60.
- 588 Sipiczki M, Kucsera J, Ulaszewski S *et al.* Hybridization Studies by Crossing and
589 Protoplast Fusion Within the Genus *Schizosaccharomyces* Lindner. *J Gen*
590 *Microbiol* 1982;**128**:1989–2000.

591 Sreeramulu G, Zhu Y, Knol W. Kombucha fermentation and its antimicrobial
592 activity. *J Agric Food Chem* 2000;**48**:2589–94.

593 Swinnen S, Thevelein JM, Nevoigt E. Genetic mapping of quantitative phenotypic
594 traits in *Saccharomyces cerevisiae*. *Fems Yeast Res* 2012;**12**:215–27.

595 Tajima F. Statistical method for testing the neutral mutation hypothesis by DNA
596 polymorphism. *Genetics* 1989;**123**:585–95.

597 Teoh AL, Heard G, Cox J. Yeast ecology of Kombucha fermentation. *Int J Food*
598 *Microbiol* 2004;**95**:119–26.

599 Teresa Avelar A, Perfeito L, Gordo I *et al*. Genome architecture is a selectable
600 trait that can be maintained by antagonistic pleiotropy. *Nat Commun*
601 2013;**4**:2235.

602 Theivendirarajah K, Chrystopher RK. Microflora and microbial activity in
603 palmyrah (*Borassus flabellifer*) palm wine in Sri Lanka. *Mircen Journal*
604 1987;**3**:23–31.

605 Vaughan Martini A. Evaluation of phylogenetic relationships among fission yeast
606 by nDNA/nDNA reassociation and conventional taxonomic criteria. *Yeast*
607 1991;**7**:73–8.

608 Vorderman AG. *Analecta Op Bromatologisch Gebied: I II*. Geneeskundig
609 Tijdschrift voor Nederlandsch-Indië, 1894:343–97.

610 Wood V, Gwilliam R, Rajandream M-A *et al*. The genome sequence of
611 *Schizosaccharomyces pombe*. *Nature* 2002;**415**:871–80.

612 Wu Q, Xu Y, Chen L. Diversity of yeast species during fermentative process
613 contributing to Chinese Maotai-flavour liquor making. *Letters in Applied*
614 *Microbiology* 2012;**55**:301–7.

615 Zanders SE, Eickbush MT, Yu JS *et al*. Genome rearrangements and pervasive
616 meiotic drive cause hybrid infertility in fission yeast. *eLife*
617 2014;**2014**:e02630.

618 Zhu YO, Siegal ML, Hall DW *et al*. Precise estimates of mutation rate and
619 spectrum in yeast. *Proc Natl Acad Sci USA* 2014;**111**:E2310–8.

620 Zimmer M, Welser F, Orlaer G *et al.* Distribution of mitochondrial introns in the
621 species *Schizosaccharomyces pombe* and the origin of the group II intron in
622 the gene encoding apocytochrome b. *Curr Genet* 1987;**12**:329–36.

623

624 The author declares that there is no conflict of interest.